

Notes

qxtr01 <qxtr01@gmail.com>

February 24, 2009

Contents

1	Symbols	2
1.1	Greek alphabet	2
2	Logic	3
2.1	Zeroth-order logic	3
2.2	First-order logic	4
3	Discrete mathematics	5
3.1	Sets	5
3.2	Binary relations	5
3.3	Binary relations over a set	6
3.4	Functions	6
3.5	Binary operations	7
3.6	Algebraic structures	7
3.7	Positional numeral systems	8
3.8	Natural numbers	8
3.9	Integer numbers	9
3.10	Real numbers	9
3.11	Polynomials	9
4	Mathematical analysis	11
4.1	Exponentiation	11
4.2	Trigonometry	11
4.3	Limits	11
5	Physics	13
5.1	Prefixes	13
6	Computer science	14
6.1	Prefixes	14

1 Symbols

1.1 Greek alphabet

A	α	alpha
B	β	beta
Γ	γ	gamma
Δ	δ	delta
E	ϵ	epsilon
Z	ζ	zeta
H	η	eta
Θ	θ	theta
I	ι	iota
K	κ	kappa
Λ	λ	lambda
M	μ	mu
N	ν	nu
Ξ	ξ	xi
O	o	omicron
Π	π	pi
P	ρ	rho
Σ	σ	sigma
T	τ	tau
Υ	υ	upsilon
Φ	ϕ	phi
X	χ	chi
Ψ	ψ	psi
Ω	ω	omega

2 Logic

2.1 Zeroth-order logic

P	\perp	\top
\perp	\perp	\top
\top	\perp	\top
P	P	$\neg P$
\perp	\perp	\top
\top	\top	\perp

P	Q	\perp	\top
\perp	\perp	\perp	\top
\perp	\top	\perp	\top
\top	\perp	\perp	\top
\top	\top	\perp	\top
P	Q	$P \wedge Q$	$P \uparrow Q$
\perp	\perp	\perp	\top
\perp	\top	\perp	\top
\top	\perp	\perp	\top
\top	\top	\top	\perp
P	Q	$P \not\rightarrow Q$	$P \rightarrow Q$
\perp	\perp	\perp	\top
\perp	\top	\perp	\top
\top	\perp	\top	\perp
\top	\top	\perp	\top
P	Q	P	$\neg P$
\perp	\perp	\perp	\top
\perp	\top	\perp	\top
\top	\perp	\top	\perp
\top	\top	\top	\perp
P	Q	$P \not\leftarrow Q$	$P \leftarrow Q$
\perp	\perp	\perp	\top
\perp	\top	\top	\perp
\top	\perp	\perp	\top
\top	\top	\perp	\top
P	Q	Q	$\neg Q$
\perp	\perp	\perp	\top
\perp	\top	\top	\perp
\top	\perp	\perp	\top
\top	\top	\top	\perp
P	Q	$P \not\leftrightarrow Q$	$P \leftrightarrow Q$
\perp	\perp	\perp	\top
\perp	\top	\top	\perp
\top	\perp	\top	\perp
\top	\top	\perp	\top
P	Q	$P \vee Q$	$P \downarrow Q$
\perp	\perp	\perp	\top
\perp	\top	\top	\perp
\top	\perp	\top	\perp
\top	\top	\top	\perp

$$\begin{aligned}
P \vee P &\iff P \\
P \vee Q &\iff Q \vee P \\
(P \vee Q) \vee R &\iff P \vee (Q \vee R)
\end{aligned}$$

$$\begin{aligned}
P \wedge P &\iff P \\
P \wedge Q &\iff Q \wedge P \\
(P \wedge Q) \wedge R &\iff P \wedge (Q \wedge R)
\end{aligned}$$

$$\begin{aligned}
\neg(P \vee Q) &\iff \neg P \wedge \neg Q \\
\neg(P \wedge Q) &\iff \neg P \vee \neg Q
\end{aligned}$$

2.2 First-order logic

$$\begin{aligned}
\forall x(P(x)) &\iff \neg \exists x(\neg P(x)) \\
\exists x(P(x)) &\iff \neg \forall x(\neg P(x)) \\
\forall x(\forall y(P(x, y))) &\iff \forall y(\forall x(P(x, y))) \\
\exists x(\exists y(P(x, y))) &\iff \exists y(\exists x(P(x, y))) \\
\forall x(P(x)) \wedge \forall x(Q(x)) &\iff \forall x(P(x) \wedge Q(x)) \\
\exists x(P(x)) \vee \exists x(Q(x)) &\iff \exists x(P(x) \vee Q(x))
\end{aligned}$$

3 Discrete mathematics

3.1 Sets

$$A \subseteq B \iff \forall x(x \in A \rightarrow x \in B)$$

$$A \subset B \iff \forall x(x \in A \rightarrow x \in B) \wedge \exists x(x \notin A \wedge x \in B)$$

$$A = B \iff \forall x(x \in A \leftrightarrow x \in B)$$

$$A = B \iff A \subseteq B \wedge B \subseteq A$$

$$A \subset B \iff A \subseteq B \wedge A \neq B$$

$$A' = \{x \in U : x \notin A\}$$

$$A \cup B = \{x \in U : x \in A \vee x \in B\}$$

$$A \cap B = \{x \in U : x \in A \wedge x \in B\}$$

$$A \setminus B = \{x \in U : x \in A \wedge x \notin B\}$$

$$A \cup A = A$$

$$A \cup B = B \cup A$$

$$(A \cup B) \cup C = A \cup (B \cup C)$$

$$A \cap A = A$$

$$A \cap B = B \cap A$$

$$(A \cap B) \cap C = A \cap (B \cap C)$$

$$(A \cup B)' = A' \cap B'$$

$$(A \cap B)' = A' \cup B'$$

3.2 Binary relations

$$R = (X, Y, G) \quad G \subseteq X \times Y$$

- total: $\forall x \in X : \exists y \in Y : (x, y) \in G$

- surjective: $\forall y \in Y : \exists x \in X : (x, y) \in G$
- functional: $\forall x \in X : \forall y_1, y_2 \in Y : (x, y_1) \in G \wedge (x, y_2) \in G \rightarrow y_1 = y_2$
- injective: $\forall y \in Y : \forall x_1, x_2 \in X : (x_1, y) \in G \wedge (x_2, y) \in G \rightarrow x_1 = x_2$
- function: total, functional
- bijective function: total, functional, injective, surjective

3.3 Binary relations over a set

$$R = (X, X, G) \quad G \subseteq X \times X$$

- reflexive: $\forall x \in X : (x, x) \in G$
- symmetric: $\forall x_1, x_2 \in X : (x_1, x_2) \in G \rightarrow (x_2, x_1) \in G$
- antisymmetric: $\forall x_1, x_2 \in X : (x_1, x_2) \in G \wedge (x_2, x_1) \in G \rightarrow x_1 = x_2$
- transitive: $\forall x_1, x_2, x_3 \in X : (x_1, x_2) \in G \wedge (x_2, x_3) \in G \rightarrow (x_1, x_3) \in G$
- equivalence relation: reflexive, symmetric, transitive
- partial order: reflexive, antisymmetric, transitive

3.4 Functions

$$f : X \rightarrow Y$$

$$x = f^{-1}(y) \mapsto y = f(x)$$

$$f^{-1} : Y \rightarrow X$$

$$y = f(x) \mapsto x = f^{-1}(y)$$

$$f : X \rightarrow Y \quad g : Y \rightarrow Z$$

$$g \circ f : X \rightarrow Z$$

$$(g \circ f)(x) = g(f(x))$$

3.5 Binary operations

$$\begin{aligned} * : X \times X &\rightarrow X \\ (x_1, x_2) &\mapsto x_1 * x_2 \end{aligned}$$

- * associative: $\forall x_1, x_2, x_3 \in X : (x_1 * x_2) * x_3 = x_1 * (x_2 * x_3)$
- * commutative: $\forall x_1, x_2 \in X : x_1 * x_2 = x_2 * x_1$
- $u_l \in X$ left identity: $\forall x \in X : u_l * x = x$
- $u_r \in X$ right identity: $\forall x \in X : x * u_r = x$
- $u \in X$ identity: $\forall x \in X : u * x = x * u = x$
- $\bar{x}_l \in X$ left inverse of $x \in X$: $\bar{x}_l * x = u$
- $\bar{x}_r \in X$ right inverse of $x \in X$: $x * \bar{x}_r = u$
- $\bar{x} \in X$ inverse of $x \in X$: $\bar{x} * x = x * \bar{x} = u$

3.6 Algebraic structures

$(X, *)$ semigroup:

1. $\forall x_1, x_2, x_3 \in X : (x_1 * x_2) * x_3 = x_1 * (x_2 * x_3)$

$(X, *)$ monoid:

1. $\forall x_1, x_2, x_3 \in X : (x_1 * x_2) * x_3 = x_1 * (x_2 * x_3)$
2. $\exists u \in X : \forall x \in X : u * x = x * u = x$

$(X, *)$ group:

1. $\forall x_1, x_2, x_3 \in X : (x_1 * x_2) * x_3 = x_1 * (x_2 * x_3)$
2. $\exists u \in X : \forall x \in X : u * x = x * u = x$
3. $\forall x \in X : \exists \bar{x} \in X : \bar{x} * x = x * \bar{x} = u$

$(X, +, \cdot)$ ring:

- $(X, +)$ Abelian group
 1. $\forall x_1, x_2, x_3 \in X : (x_1 + x_2) + x_3 = x_1 + (x_2 + x_3)$
 2. $\exists 0 \in X : \forall x \in X : 0 + x = x + 0 = x$
 3. $\forall x \in X : \exists (-x) \in X : (-x) + x = x + (-x) = 0$
 4. $\forall x_1, x_2 \in X : x_1 + x_2 = x_2 + x_1$
- (R, \cdot) monoid
 1. $\forall x_1, x_2, x_3 \in X : (x_1 \cdot x_2) \cdot x_3 = x_1 \cdot (x_2 \cdot x_3)$
 2. $\exists 1 \in X : \forall x \in X : 1 \cdot x = x \cdot 1 = x$
- $\forall x_1, x_2, x_3 \in X : (x_1 + x_2) \cdot x_3 = (x_1 \cdot x_3) + (x_2 \cdot x_3)$

- $\forall x_1, x_2, x_3 \in X : x_1 \cdot (x_2 + x_3) = (x_1 \cdot x_2) + (x_1 \cdot x_3)$

$(X, +, \cdot)$ field:

- $(X, +)$ Abelian group

1. $\forall x_1, x_2, x_3 \in X : (x_1 + x_2) + x_3 = x_1 + (x_2 + x_3)$
2. $\exists 0 \in X : \forall x \in X : 0 + x = x + 0 = x$
3. $\forall x \in X : \exists (-x) \in X : (-x) + x = x + (-x) = 0$
4. $\forall x_1, x_2 \in X : x_1 + x_2 = x_2 + x_1$

- (R, \cdot) Abelian monoid

1. $\forall x_1, x_2, x_3 \in X : (x_1 \cdot x_2) \cdot x_3 = x_1 \cdot (x_2 \cdot x_3)$
2. $\exists 1 \in X : \forall x \in X : 1 \cdot x = x \cdot 1 = x$
3. $\forall x_1, x_2 \in X : x_1 \cdot x_2 = x_2 \cdot x_1$

- $\forall x_1, x_2, x_3 \in X : (x_1 + x_2) \cdot x_3 = (x_1 \cdot x_3) + (x_2 \cdot x_3)$

- $\forall x_1, x_2, x_3 \in X : x_1 \cdot (x_2 + x_3) = (x_1 \cdot x_2) + (x_1 \cdot x_3)$

- $\forall x \neq 0 : \exists x^{-1} : x \cdot x^{-1} = x^{-1} \cdot x = 1$

3.7 Positional numeral systems

$$b \in \mathbb{N} \setminus \{0, 1\}$$

$$a_0, \dots, a_m \in \mathbb{R}$$

$$c_1, \dots, c_n \in \mathbb{R}$$

$$(a_m \cdots a_0.c_1 \cdots c_n)_b = \sum_{i=0}^m a_i b^i + \sum_{i=1}^n c_i b^{-i}$$

3.8 Natural numbers

$$n \in \mathbb{N}$$

$$\sum_{i=0}^n i = \frac{n(n+1)}{2}$$

$$n \in \mathbb{N}$$

$$n! = \begin{cases} 1 & n = 0 \\ n(n-1)! & n \geq 1 \end{cases}$$

3.9 Integer numbers

$$a \in \mathbb{Z} \setminus \{0\} \quad b \in \mathbb{Z}$$
$$a \mid b \iff \exists h \in \mathbb{Z} : b = ha$$

$$\text{gcd} : \mathbb{Z}^2 \setminus \{(0, 0)\} \rightarrow \mathbb{N} \setminus \{0\}$$
$$(z_1, z_2) \mapsto \text{gcd}(z_1, z_2)$$

$$\text{lcm} : \mathbb{Z} \setminus \{0\} \times \mathbb{Z} \setminus \{0\} \rightarrow \mathbb{N} \setminus \{0\}$$
$$(z_1, z_2) \mapsto \text{lcm}(z_1, z_2)$$

3.10 Real numbers

$$\sqrt{2} = 1.4142135623\dots$$

$$\sqrt{3} = 1.7320508075\dots$$

$$\sqrt{5} = 2.2360679774\dots$$

$$e = 2.7182818284\dots$$

$$\pi = 3.1415926535\dots$$

3.11 Polynomials

$$p : \mathbb{R} \rightarrow \mathbb{R}$$
$$p(x) = \sum_{i=0}^n a_i x^i \quad n \in \mathbb{N}$$

$$\text{deg} : \mathbb{R}[x] \setminus \{0_{\mathbb{R}[x]}\} \rightarrow \mathbb{N}$$

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x - y)^2 = x^2 - 2xy + y^2$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$$

$$(x + y)(x - y) = x^2 - y^2$$

$$(x + y)(x^2 - xy + y^2) = x^3 + y^3$$

$$(x + y)(x^2 + xy + y^2) = x^3 - y^3$$

$$ax^2 + bx + c = 0 \quad a \neq 0$$
$$b^2 - 4ac \geq 0 \implies x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$
$$ax^2 + bx + c = a(x - x_1)(x - x_2)$$

4 Mathematical analysis

4.1 Exponentiation

$$a \in \mathbb{R} \quad n \in \mathbb{N}$$
$$a^n = \begin{cases} 1 & n = 0 \\ a \cdot a^{n-1} & n \geq 1 \end{cases}$$

$$a^{m+n} = a^m \cdot a^n$$
$$(a^m)^n = a^{m \cdot n}$$
$$(a \cdot b)^n = a^n \cdot b^n$$

4.2 Trigonometry

$$\sin(x \pm y) = \sin(x) \cos(y) \pm \cos(x) \sin(y)$$
$$\cos(x \pm y) = \cos(x) \cos(y) \mp \sin(x) \sin(y)$$

$$\sin(2x) = 2 \sin(x) \cos(x)$$
$$\cos(2x) = 2 \cos^2(x) - 1$$

4.3 Limits

$$x_0 \in \mathbb{R} \quad \delta \in \{x \in \mathbb{R} : x > 0\}$$
$$I_\delta(x_0) = (x_0 - \delta, x_0 + \delta)$$
$$I_\delta^-(x_0) = (x_0 - \delta, x_0]$$
$$I_\delta^+(x_0) = [x_0, x_0 + \delta)$$

$$\delta \in \mathbb{R}$$
$$I_\delta(+\infty) = (\delta, +\infty)$$
$$I_\delta(-\infty) = (-\infty, \delta)$$

$$x_0 \in \bar{\mathbb{R}} \quad A \subseteq \mathbb{R}$$

$$x_0 \in D(A) \iff \forall I_\delta(x_0) \in I(x_0) : A \cap I_\delta(x_0) \setminus \{x_0\} \neq \emptyset$$

$$f : D_f \subseteq \mathbb{R} \rightarrow \mathbb{R} \quad x_0 \in D(D_f) \quad l \in \bar{\mathbb{R}}$$

$$\lim_{x \rightarrow x_0} f(x) = l$$

$$\forall I_\epsilon(l) \in I(l) : \exists I_\delta(x_0) \in I(x_0) : \forall x \in I_\delta(x_0) \setminus \{x_0\} : f(x) \in I_\epsilon(l)$$

$$f : D_f \subseteq \mathbb{R} \rightarrow \mathbb{R} \quad x_0 \in D(D_f \cap I_{x_0}(\pm\infty)) \quad l \in \bar{\mathbb{R}}$$

$$\lim_{x \rightarrow x_0^\pm} f(x) = l$$

$$\forall I_\epsilon(l) \in I(l) : \exists I_\delta^\pm(x_0) \in I^\pm(x_0) : \forall x \in I_\delta^\pm(x_0) \setminus \{x_0\} : f(x) \in I_\epsilon(l)$$

5 Physics

5.1 Prefixes

10^{18}	exa	E
10^{15}	peta	P
10^{12}	tera	T
10^9	giga	G
10^6	mega	M
10^3	kilo	k
10^2	hecto	h
10^1	deca	da
10^0		
10^{-1}	deci	d
10^{-2}	centi	c
10^{-3}	milli	m
10^{-6}	micro	μ
10^{-9}	nano	n
10^{-12}	pico	p
10^{-15}	femto	f
10^{-18}	atto	a

6 Computer science

6.1 Prefixes

2^{60}	exbi	Ei
2^{50}	pebi	Pi
2^{40}	tebi	Ti
2^{30}	gibi	Gi
2^{20}	mebi	Mi
2^{10}	kibi	Ki